



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



On a non-homogeneous eigenvalue problem involving a potential: An Orlicz–Sobolev space setting

Mihai Mihăilescu^{a,b,*}, Vicențiu Rădulescu^{a,c,d}, Dušan Repovš^e

^a Department of Mathematics, University of Craiova, 200585 Craiova, Romania

^b Department of Mathematics, Central European University, 1051 Budapest, Hungary

^c Institute of Mathematics “Simion Stoilow” of the Romanian Academy, PO Box 1-764, 014700 Bucharest, Romania

^d Institute of Mathematics, Physics and Mechanics, POB 2964, Ljubljana, Slovenia 1001

^e Faculty of Mathematics and Physics, and Faculty of Education, University of Ljubljana, POB 2964, Ljubljana, Slovenia 1001

Received 27 March 2009

Available online 3 June 2009

Abstract

In this paper we study a non-homogeneous eigenvalue problem involving variable growth conditions and a potential V . The problem is analyzed in the context of Orlicz–Sobolev spaces. Connected with this problem we also study the optimization problem for the particular eigenvalue given by the infimum of the Rayleigh quotient associated to the problem with respect to the potential V when V lies in a bounded, closed and convex subset of a certain variable exponent Lebesgue space.

© 2009 Elsevier Masson SAS. All rights reserved.

Résumé

Dans cet article on étudie un problème non homogène à valeurs propres avec exposant variable et potentiel V . Ce problème est analysé dans les espaces d'Orlicz–Sobolev. On étudie également le problème d'optimisation dans le cas particulier où la valeur propre s'obtient par minimisation du quotient de Rayleigh associé au potentiel V , quand V appartient à un ensemble borné, fermé et convexe d'un espace de Lebesgue à exposant variable.

© 2009 Elsevier Masson SAS. All rights reserved.

MSC: 35D05; 35J60; 35J70; 58E05; 68T40; 76A02

Keywords: Eigenvalue problem; Orlicz–Sobolev space; Variable exponent Lebesgue space; Optimization problem

1. Introduction and preliminary results

Let Ω be a bounded domain in \mathbb{R}^N ($N \geq 3$) with smooth boundary $\partial\Omega$. Assume that $a_i : (0, \infty) \rightarrow \mathbb{R}$, $i = 1, 2$, are two functions such that the mappings $\varphi_i : \mathbb{R} \rightarrow \mathbb{R}$, $i = 1, 2$, defined by:

* Corresponding author at: Department of Mathematics, University of Craiova, 200585 Craiova, Romania.

E-mail addresses: mmihailles@yahoo.com (M. Mihăilescu), vicentiu.radulescu@imar.ro (V. Rădulescu), dusan.repovs@guest.arnes.si (D. Repovš).

$$\varphi_i(t) = \begin{cases} a_i(|t|)t, & \text{for } t \neq 0, \\ 0, & \text{for } t = 0, \end{cases}$$

are odd, increasing homeomorphisms from \mathbb{R} onto \mathbb{R} , λ is a real number, $V(x)$ is a potential and $q_1, q_2, m : \overline{\Omega} \rightarrow (1, \infty)$ are continuous functions. We analyze the eigenvalue problem:

$$\begin{cases} -\operatorname{div}((a_1(|\nabla u|) + a_2(|\nabla u|))\nabla u) + V(x)|u|^{m(x)-2}u = \lambda(|u|^{q_1(x)-2} + |u|^{q_2(x)-2})u, & \text{if } x \in \Omega \\ u = 0, & \text{if } x \in \partial\Omega. \end{cases} \quad (1)$$

The interest in analyzing this kind of problems is motivated by some recent advances in the study of eigenvalue problems involving non-homogeneous operators in the divergence form. We refer especially to the results in [13, 18,20,12,21–23]. Problem (1) can be placed in the context of the above results since in the particular case when $q_1(x) = q_2(x) = q(x)$ for any $x \in \overline{\Omega}$ and $V \equiv 0$ in Ω it was studied in [21]. The form of problem (1) becomes a natural extension of the problem studied in [21] with the presence of the potential V in the left-hand side of the equation and by considering that in the right-hand side we can have $q_1 \neq q_2$ on $\overline{\Omega}$.

In order to go further we introduce the functional space setting where problem (1) will be discussed. In this context we notice that the operator in the divergence form is not homogeneous and thus, we introduce an Orlicz–Sobolev space setting for problems of this type. On the other hand, the presence of the continuous functions m, q_1 and q_2 as exponents appeals to a suitable variable exponent Lebesgue space setting. In the following, we give a brief description of the Orlicz–Sobolev spaces and of the variable exponent Lebesgue spaces.

We start by recalling some basic facts about Orlicz spaces. For more details we refer to the books by D.R. Adams and L.I. Hedberg [2], R. Adams [3] and M.M. Rao and Z.D. Ren [25] and the papers by Ph. Clément et al. [6,7], M. García-Huidobro et al. [14] and J.P. Gossez [15].

For $\varphi_i : \mathbb{R} \rightarrow \mathbb{R}, i = 1, 2$, which are odd, increasing homeomorphisms from \mathbb{R} onto \mathbb{R} , we define

$$\Phi_i(t) = \int_0^t \varphi_i(s) ds, \quad (\Phi_i)^*(t) = \int_0^t (\varphi_i)^{-1}(s) ds, \quad \text{for all } t \in \mathbb{R}, i = 1, 2.$$

We observe that $\Phi_i, i = 1, 2$, are *Young functions*, that is, $\Phi_i(0) = 0, \Phi_i$ are convex, and $\lim_{x \rightarrow \infty} \Phi_i(x) = +\infty$. Furthermore, since $\Phi_i(x) = 0$ if and only if $x = 0, \lim_{x \rightarrow 0} \Phi_i(x)/x = 0$, and $\lim_{x \rightarrow \infty} \Phi_i(x)/x = +\infty$, then Φ_i are called *N-functions*. The functions $(\Phi_i)^*, i = 1, 2$, are called the *complementary functions* of $\Phi_i, i = 1, 2$, and they satisfy:

$$(\Phi_i)^*(t) = \sup\{st - \Phi_i(s); s \geq 0\}, \quad \text{for all } t \geq 0.$$

We also observe that $(\Phi_i)^*, i = 1, 2$, are also *N-functions* and Young's inequality holds true

$$st \leq \Phi_i(s) + (\Phi_i)^*(t), \quad \text{for all } s, t \geq 0.$$

The Orlicz spaces $L_{\Phi_i}(\Omega), i = 1, 2$, defined by the *N-functions* Φ_i (see [2,3,6]) are the spaces of measurable functions $u : \Omega \rightarrow \mathbb{R}$ such that

$$\|u\|_{L_{\Phi_i}} := \sup \left\{ \int_{\Omega} uv dx; \int_{\Omega} (\Phi_i)^*(|g|) dx \leq 1 \right\} < \infty.$$

Then $(L_{\Phi_i}(\Omega), \|\cdot\|_{L_{\Phi_i}}), i = 1, 2$, are Banach spaces whose norm is equivalent to the Luxemburg norm

$$\|u\|_{\Phi_i} := \inf \left\{ k > 0; \int_{\Omega} \Phi_i \left(\frac{u(x)}{k} \right) dx \leq 1 \right\}.$$

For Orlicz spaces Hölder's inequality reads as follows (see [25, Inequality 4, p. 79]):

$$\int_{\Omega} uv dx \leq 2\|u\|_{L_{\Phi_i}} \|v\|_{L_{(\Phi_i)^*}} \quad \text{for all } u \in L_{\Phi_i}(\Omega) \text{ and } v \in L_{(\Phi_i)^*}(\Omega), i = 1, 2.$$

Next, we introduce the Orlicz–Sobolev spaces. We denote by $W^1L_{\Phi_i}(\Omega)$, $i = 1, 2$, the Orlicz–Sobolev spaces defined by:

$$W^1L_{\Phi_i}(\Omega) := \left\{ u \in L_{\Phi_i}(\Omega); \frac{\partial u}{\partial x_i} \in L_{\Phi_i}(\Omega), i = 1, \dots, N \right\}.$$

These are Banach spaces with respect to the norms

$$\|u\|_{1, \Phi_i} := \|u\|_{\Phi_i} + \|\nabla u\|_{\Phi_i}, \quad i = 1, 2.$$

We also define the Orlicz–Sobolev spaces $W_0^1L_{\Phi_i}(\Omega)$, $i = 1, 2$, as the closure of $C_0^\infty(\Omega)$ in $W^1L_{\Phi_i}(\Omega)$. By Lemma 5.7 in [15] we obtain that on $W_0^1L_{\Phi_i}(\Omega)$, $i = 1, 2$, we may consider some equivalent norms

$$\|u\|_i := \|\nabla u\|_{\Phi_i}.$$

For an easier manipulation of the spaces defined above, we define:

$$(\varphi_i)_0 := \inf_{t>0} \frac{t\varphi_i(t)}{\Phi_i(t)} \quad \text{and} \quad (\varphi_i)^0 := \sup_{t>0} \frac{t\varphi_i(t)}{\Phi_i(t)}, \quad i \in \{1, 2\}.$$

In this paper we assume that for each $i \in \{1, 2\}$ we have:

$$1 < (\varphi_i)_0 \leq \frac{t\varphi_i(t)}{\Phi_i(t)} \leq (\varphi_i)^0 < \infty, \quad \forall t \geq 0. \tag{2}$$

The above relation implies that each Φ_i , $i \in \{1, 2\}$, satisfies the Δ_2 -condition, i.e.

$$\Phi_i(2t) \leq K\Phi_i(t), \quad \forall t \geq 0, \tag{3}$$

where K is a positive constant (see [22, Proposition 2.3]).

On the other hand, the following relations hold true

$$\|u\|_i^{(\varphi_i)_0} \leq \int_{\Omega} \Phi_i(|\nabla u|) dx \leq \|u\|_i^{(\varphi_i)^0}, \quad \forall u \in W_0^1L_{\Phi_i}(\Omega) \text{ with } \|u\|_i < 1, i = 1, 2, \tag{4}$$

$$\|u\|_i^{(\varphi_i)^0} \leq \int_{\Omega} \Phi_i(|\nabla u|) dx \leq \|u\|_i^{(\varphi_i)_0}, \quad \forall u \in W_0^1L_{\Phi_i}(\Omega) \text{ with } \|u\|_i > 1, i = 1, 2 \tag{5}$$

(see, e.g. [21, Lemma 1]).

Furthermore, in this paper we assume that for each $i \in \{1, 2\}$ the function Φ_i satisfies the following condition:

$$\text{the function } [0, \infty) \ni t \rightarrow \Phi_i(\sqrt{t}) \text{ is convex.} \tag{6}$$

Conditions (3) and (6) assure that for each $i \in \{1, 2\}$ the Orlicz spaces $L_{\Phi_i}(\Omega)$ are uniformly convex spaces and thus, reflexive Banach spaces (see [22, Proposition 2.2]). That fact implies that also the Orlicz–Sobolev spaces $W_0^1L_{\Phi_i}(\Omega)$, $i \in \{1, 2\}$, are reflexive Banach spaces.

Remark 1. We point out certain examples of functions $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ which are odd, increasing homeomorphisms from \mathbb{R} onto \mathbb{R} and satisfy conditions (2) and (6). For more details the reader can consult [7, Examples 1–3, p. 243].

1) Let

$$\varphi(t) = p|t|^{p-2}t, \quad \forall t \in \mathbb{R},$$

with $p > 1$. For this function it can be proved that

$$(\varphi)_0 = (\varphi)^0 = p.$$

Furthermore, in this particular case the corresponding Orlicz space $L_{\Phi}(\Omega)$ is the classical Lebesgue space $L^p(\Omega)$ while the Orlicz–Sobolev space $W_0^1L_{\Phi}(\Omega)$ is the classical Sobolev space $W_0^{1,p}(\Omega)$. We will use the classical notations to denote the Orlicz–Sobolev spaces in this particular case.

2) Consider

$$\varphi(t) = \log(1 + |t|^s)|t|^{p-2}t, \quad \forall t \in \mathbb{R},$$

with $p, s > 1$. In this case it can be proved that

$$(\varphi)_0 = p, \quad (\varphi)^0 = p + s.$$

3) Let

$$\varphi(t) = \frac{|t|^{p-2}t}{\log(1 + |t|)}, \quad \text{if } t \neq 0, \varphi(0) = 0,$$

with $p > 2$. In this case we have

$$(\varphi)_0 = p - 1, \quad (\varphi)^0 = p.$$

Next, we recall some background facts concerning the variable exponent Lebesgue spaces. For more details we refer to the book by Musielak [24] and the papers by Edmunds et al. [8–10], Kováčik and Rákosník [16], Mihăilescu and Rădulescu [17], and Samko and Vakulov [26].

Set

$$C_+(\overline{\Omega}) = \{h; h \in C(\overline{\Omega}), h(x) > 1 \text{ for all } x \in \overline{\Omega}\}.$$

For any $h \in C_+(\overline{\Omega})$ we define:

$$h^+ = \sup_{x \in \Omega} h(x) \quad \text{and} \quad h^- = \inf_{x \in \Omega} h(x).$$

For any $q(x) \in C_+(\overline{\Omega})$ we define the variable exponent Lebesgue space $L^{q(x)}(\Omega)$ (see [16]). On $L^{q(x)}(\Omega)$ we define the Luxemburg norm by the formula

$$|u|_{q(x)} = \inf \left\{ \mu > 0; \int_{\Omega} \left| \frac{u(x)}{\mu} \right|^{q(x)} dx \leq 1 \right\}.$$

We remember that the variable exponent Lebesgue spaces are separable and reflexive Banach spaces. If $0 < |\Omega| < \infty$ and q_1, q_2 are variable exponents so that $q_1(x) \leq q_2(x)$ almost everywhere in Ω then there exists the continuous embedding $L^{q_2(x)}(\Omega) \hookrightarrow L^{q_1(x)}(\Omega)$.

Let $L^{p'(x)}(\Omega)$ denote the conjugate space of $L^{p(x)}(\Omega)$, where $1/p(x) + 1/p'(x) = 1$. For any $u \in L^{p(x)}(\Omega)$ and $v \in L^{p'(x)}(\Omega)$ the Hölder type inequality

$$\left| \int_{\Omega} uv \, dx \right| \leq \left(\frac{1}{p^-} + \frac{1}{p'^-} \right) |u|_{p(x)} |v|_{p'(x)}, \tag{7}$$

holds true.

If $(u_n), u \in L^{q(x)}(\Omega)$ then the following relations hold true:

$$|u|_{q(x)} > 1 \quad \Rightarrow \quad |u|_{q(x)}^{q^-} \leq \int_{\Omega} |u|^{q(x)} dx \leq |u|_{q(x)}^{q^+}, \tag{8}$$

$$|u|_{q(x)} < 1 \quad \Rightarrow \quad |u|_{q(x)}^{q^+} \leq \int_{\Omega} |u|^{q(x)} dx \leq |u|_{q(x)}^{q^-}, \tag{9}$$

$$|u_n - u|_{q(x)} \rightarrow 0 \quad \Leftrightarrow \quad \int_{\Omega} |u_n - u|^{q(x)} dx \rightarrow 0. \tag{10}$$

Now we can turn back to problem (1). We will study problem (1) when $q_1, q_2, m : \overline{\Omega} \rightarrow (1, \infty)$ are continuous functions satisfying the following assumptions:

$$1 < (\varphi_2)_0 \leq (\varphi_2)^0 < q_2^- \leq q_2^+ \leq m^- \leq m^+ \leq q_1^- \leq q_1^+ < (\varphi_1)_0 \leq (\varphi_1)^0 < N, \tag{11}$$

$$q_1^+ < [(\varphi_2)_0]^* := \frac{N(\varphi_2)_0}{N - (\varphi_2)_0}, \quad \forall x \in \overline{\Omega}, \tag{12}$$

and the potential $V : \Omega \rightarrow \mathbb{R}$ satisfies

$$V \in L^{r(x)}(\Omega), \quad \text{with } r(x) \in C(\overline{\Omega}) \quad \text{and} \quad r(x) > \frac{N}{m}, \quad \forall x \in \overline{\Omega}. \tag{13}$$

Condition (11) which describes the competition between the growth rates involved in Eq. (1), actually, assures a balance between them and thus, it represents the *key* of the present study. Such a balance is essential since we are working on a non-homogeneous (eigenvalue) problem for which a minimization technique based on the Lagrange Multiplier Theorem cannot be applied in order to find (principal) eigenvalues (unlike the case offered by the homogeneous operators). Thus, in the case of nonlinear non-homogeneous eigenvalue problems the classical theory used in the homogeneous case does not work entirely, but some of its ideas can still be useful and some particular results can still be obtained in some aspects while in other aspects entirely new phenomena can occur. To focus on our case, condition (11) together with conditions (12) and (13) imply:

$$\lim_{\|u\|_1 \rightarrow 0} \frac{\int_{\Omega} \Phi_1(|\nabla u|) dx + \int_{\Omega} \Phi_2(|\nabla u|) dx + \int_{\Omega} \frac{V(x)}{m(x)} |u|^{m(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u|^{q_2(x)} dx} = \infty,$$

and

$$\lim_{\|u\|_1 \rightarrow \infty} \frac{\int_{\Omega} \Phi_1(|\nabla u|) dx + \int_{\Omega} \Phi_2(|\nabla u|) dx + \int_{\Omega} \frac{V(x)}{m(x)} |u|^{m(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u|^{q_2(x)} dx} = \infty.$$

In other words, the absence of homogeneity is balanced by the behavior (actually, the blow-up) of the Rayleigh quotient associated to problem (1) in the origin and at infinity. The consequences of the above remarks is that the infimum of the Rayleigh quotient associated to problem (1) is a real number, i.e.

$$\inf_{u \in W_0^1 L_{\Phi_1}(\Omega) \setminus \{0\}} \frac{\int_{\Omega} \Phi_1(|\nabla u|) dx + \int_{\Omega} \Phi_2(|\nabla u|) dx + \int_{\Omega} \frac{V(x)}{m(x)} |u|^{m(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u|^{q_2(x)} dx} \in \mathbb{R}, \tag{14}$$

and it will be attained for a function $u_0 \in W_0^{1,p_1(x)}(\Omega) \setminus \{0\}$. Moreover, the value in (14) represents an eigenvalue of problem (1) with the corresponding eigenfunction u_0 . However, at this stage we cannot say if the eigenvalue described above is the lowest eigenvalue of problem (1) or not, even if we are able to show that any λ small enough is not an eigenvalue of (1). For the moment this rests an open question. On the other hand, we can prove that any λ superior to the value given by relation (14) is also an eigenvalue of problem (1). Thus, we conclude that problem (1) possesses a continuous family of eigenvalues.

Related with the above ideas we will also discuss the *optimization* of the eigenvalues described by relation (14) with respect to the potential V , providing that V belongs to a bounded, closed and convex subset of $L^{r(x)}(\Omega)$ (where $r(x)$ is given by relation (13)). By optimization we understand the existence of some potentials V_* and V^* such that the eigenvalue described in relation (14) is minimal or maximal with respect to the set where V lies. The results that we will obtain in the context of optimization of eigenvalues are motivated by the above advances in this field in the case of homogeneous (linear or nonlinear) eigenvalue problems. We refer mainly to the studies in Ashbaugh and Harrell [1], Egnell [11] and Bonder and Del Pezzo [4] where different optimization problems of the principal eigenvalue of some homogeneous operators were studied.

2. The main results

By relation (11) it follows that $W_0^1 L_{\Phi_1}(\Omega)$ is continuously embedded in $W_0^1 L_{\Phi_2}(\Omega)$ (see, e.g. [21, Lemma 2]). Thus, problem (1) will be analyzed in the space $W_0^1 L_{\Phi_1}(\Omega)$.

We say that $\lambda \in \mathbb{R}$ is an *eigenvalue* of problem (1) if there exists $u \in W_0^1 L_{\Phi_1}(\Omega) \setminus \{0\}$ such that

$$\int_{\Omega} (a_1(|\nabla u|) + a_2(|\nabla u|)) \nabla u \nabla v dx + \int_{\Omega} V(x) |u|^{m(x)-2} uv dx - \lambda \int_{\Omega} (|u|^{q_1(x)-2} + |u|^{q_2(x)-2}) uv dx = 0,$$

for all $v \in W_0^1 L_{\Phi_1}(\Omega)$. We point out that if λ is an eigenvalue of problem (1) then the corresponding *eigenfunction* $u \in W_0^1 L_{\Phi_1}(\Omega) \setminus \{0\}$ is a *weak solution* of problem (1).

For each potential $V \in L^{r(x)}(\Omega)$ we define:

$$A(V) := \inf_{u \in W_0^1 L_{\Phi_1}(\Omega) \setminus \{0\}} \frac{\int_{\Omega} \Phi_1(|\nabla u|) dx + \int_{\Omega} \Phi_2(|\nabla u|) dx + \int_{\Omega} \frac{V(x)}{m(x)} |u|^{m(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u|^{q_2(x)} dx},$$

and

$$B(V) := \inf_{u \in W_0^1 L_{\Phi_1}(\Omega) \setminus \{0\}} \frac{\int_{\Omega} a_1(|\nabla u|) |\nabla u|^2 dx + \int_{\Omega} a_2(|\nabla u|) |\nabla u|^2 dx + \int_{\Omega} V(x) |u|^{m(x)} dx}{\int_{\Omega} |u|^{q_1(x)} dx + \int_{\Omega} |u|^{q_2(x)} dx}.$$

Thus, we can define two functions $A, B : L^{r(x)}(\Omega) \rightarrow \mathbb{R}$.

The first result of this paper is given by the following theorem.

Theorem 2.1. *Assume that conditions (11), (12) and (13) are fulfilled. Then $A(V)$ is an eigenvalue of problem (1). Moreover, there exists $u_V \in W_0^1 L_{\Phi_1}(\Omega) \setminus \{0\}$ an eigenfunction corresponding to the eigenvalue $A(V)$ such that*

$$A(V) = \frac{\int_{\Omega} \Phi_1(|\nabla u_V|) dx + \int_{\Omega} \Phi_2(|\nabla u_V|) dx + \int_{\Omega} \frac{V(x)}{m(x)} |u_V|^{m(x)} dx}{\int_{\Omega} \frac{1}{q_1(x)} |u_V|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u_V|^{q_2(x)} dx}.$$

Furthermore, $B(V) \leq A(V)$, each $\lambda \in (A(V), \infty)$ is an eigenvalue of problem (1), while each $\lambda \in (-\infty, B(V))$ is not an eigenvalue of problem (1).

Next, we will show that on each convex, bounded and closed subset of $L^{r(x)}(\Omega)$ the function A defined above is bounded from below and attains its minimum. The result is the following:

Theorem 2.2. *Assume that conditions (11), (12) and (13) are fulfilled. Assume that S is a convex, bounded and closed subset of $L^{r(x)}(\Omega)$. Then there exists $V_{\star} \in S$ which minimizes $A(V)$ on S , i.e.*

$$A(V_{\star}) = \inf_{V \in S} A(V).$$

Finally, we will focus our attention on the particular case when the set S from Theorem 2.2 is a ball in $L^{r(x)}(\Omega)$. Thus, we will denote each closed ball centered in the origin of radius R from $L^{r(x)}(\Omega)$ by $\bar{B}_R(0)$, i.e.

$$\bar{B}_R(0) := \{u \in L^{r(x)}(\Omega); |u|_{r(x)} \leq R\}.$$

By Theorem 2.2 we can define the function $A_{\star} : [0, \infty) \rightarrow \mathbb{R}$ by:

$$A_{\star}(R) = \min_{V \in \bar{B}_R(0)} A(V).$$

Our result on the function A_{\star} is given by the following theorem:

Theorem 2.3.

- (a) *The function A_{\star} is not constant and decreases monotonically.*
- (b) *The function A_{\star} is continuous.*

On the other hand, we point out that similar results as those of Theorems 2.2 and 2.3 can be obtained if we notice that on each convex, bounded and closed subset of $L^{r(x)}(\Omega)$ the function A defined in Theorem 2.1 is also bounded from above and attains its maximum. It is also easy to remark that we can define a function $A^{\star} : [0, \infty) \rightarrow \mathbb{R}$ by:

$$A^{\star}(R) = \max_{V \in \bar{B}_R(0)} A(V),$$

which has similar properties as A_{\star} .

3. Proof of Theorem 2.1

Let X denote the generalized Sobolev space $W_0^1 L_{\Phi_1}(\Omega)$. Relation (11) and similar arguments as those used in [21, Lemma 1] combined with [3, Lemma 8.12(b)] and with the Rellich–Kondrachov theorem we deduce that

$$W_0^1 L_{\Phi_1}(\Omega) \subset W_0^1 L_{\Phi_2}(\Omega) \subset W_0^{1,(\varphi_2)_0}(\Omega) \hookrightarrow L^{q_1^+}(\Omega) \subset L^{q_1(x)}(\Omega) \subset L^{m(x)}(\Omega) \subset L^{q_2(x)}(\Omega), \quad (15)$$

where we denoted by \subset a *continuous* embedding while by \hookrightarrow we denoted a *compact* embedding.

Define the functionals $J_V, I : X \rightarrow \mathbb{R}$ by:

$$J_V(u) = \int_{\Omega} \Phi_1(|\nabla u|) dx + \int_{\Omega} \Phi_2(|\nabla u|) dx + \int_{\Omega} \frac{V(x)}{m(x)} |u|^{m(x)} dx,$$

$$I(u) = \int_{\Omega} \frac{1}{q_1(x)} |u|^{q_1(x)} dx + \int_{\Omega} \frac{1}{q_2(x)} |u|^{q_2(x)} dx.$$

Relation (15) assures that the functionals defined above are well defined. We notice that for any V satisfying condition (13) we have:

$$J_V(u) = J_0(u) + \int_{\Omega} \frac{V(x)}{m(x)} |u|^{m(x)} dx, \quad \forall u \in X,$$

where J_0 is obtained in the case when $V = 0$ in Ω .

Standard arguments imply that $J_V, I \in C^1(X, \mathbb{R})$ and for all $u, v \in X$,

$$\langle J'_V(u), v \rangle = \int_{\Omega} (a_1(|\nabla u|) + a_2(|\nabla u|)) \nabla u \nabla v dx + \int_{\Omega} V(x) |u|^{m(x)-2} uv dx,$$

$$\langle I'(u), v \rangle = \int_{\Omega} |u|^{q_1(x)-2} uv dx + \int_{\Omega} |u|^{q_2(x)-2} uv dx.$$

In order to prove Theorem 2.1 we first establish some auxiliary results.

Lemma 3.1. *Assume conditions (11), (12) and (13) are fulfilled. Then for each $\epsilon > 0$ there exists $C_{\epsilon} > 0$ such that*

$$\left| \int_{\Omega} \frac{V(x)}{m(x)} |u|^{m(x)} dx \right| \leq \epsilon \int_{\Omega} (\Phi_1(|\nabla u|) + \Phi_2(|\nabla u|)) dx + C_{\epsilon} |V|_{r(x)} \int_{\Omega} (|u|^{m^-} + |u|^{m^+}) dx,$$

for all $u \in X$.

Proof. First, we point out that since $r(x) > r^-$ on $\overline{\Omega}$ it follows that $L^{r(x)}(\Omega) \subset L^{r^-}(\Omega)$. On the other hand, since $r(x) > \frac{N}{m^-}$ for each $x \in \overline{\Omega}$ it follows that $r^- > \frac{N}{m^-}$. Thus, we infer that $V \in L^{r^-}(\Omega)$ and $r^- > \frac{N}{m^-}$.

Now, let $\epsilon > 0$ be fixed. We claim that there exists $D_{\epsilon} > 0$ such that

$$\int_{\Omega} |V(x)| \cdot |u|^{m^-} dx \leq \epsilon \int_{\Omega} |\nabla u|^{m^-} dx + D_{\epsilon} |V|_{r^-} \int_{\Omega} |u|^{m^-} dx, \quad \forall u \in W_0^{1,m^-}(\Omega). \quad (16)$$

In order to establish (16) we show first that for each $s \in (1, \frac{Nm^-}{N-m^-})$ there exists $D'_{\epsilon} > 0$ such that

$$|v|_s \leq \epsilon \|\nabla v\|_{m^-} + D'_{\epsilon} |v|_{m^-}, \quad \forall v \in W_0^{1,m^-}(\Omega). \quad (17)$$

Indeed, assume by contradiction that relation (17) does not hold true for each $\epsilon > 0$. Then there exists $\epsilon_0 > 0$ and a sequence $(v_n) \subset W_0^{1,m^-}(\Omega)$ such that $|v_n|_s = 1$ and

$$\epsilon_0 \|\nabla v_n\|_{m^-} + n |v_n|_{m^-} < 1, \quad \forall n.$$

Then it is clear that (v_n) is bounded in $W_0^{1,m^-}(\Omega)$ and $|v_n|_{m^-} \rightarrow 0$. Thus, we deduce that passing eventually to a subsequence we can assume that v_n converges weakly to a function v in $W_0^{1,m^-}(\Omega)$ and actually $v = 0$. Since $s \in (1, \frac{Nm^-}{N-m^-})$ it follows by the Rellich–Kondrachov theorem that $W_0^{1,m^-}(\Omega)$ is compactly embedded in $L^s(\Omega)$ and thus v_n converges to 0 in $L^s(\Omega)$. On the other hand, since $|v_n|_s = 1$ for each n we deduce that $|v|_s = 1$ and that is a contradiction. We obtained that relation (17) holds true.

Next, we point out that since $r^- > \frac{N}{m^-}$ then $m^- \cdot r^{-'} < \frac{Nm^-}{N-m^-}$, where $r^{-'} = \frac{r^-}{r^- - 1}$. Thus, by Hölder’s inequality we have:

$$\int_{\Omega} |V(x)| \cdot |u|^{m^-} dx \leq |V|_{r^-} \cdot |u|_{m^- \cdot r^{-'}}^{m^-}, \quad \forall u \in W_0^{1,m^-}(\Omega).$$

Combining the last inequality with relation (17) we infer that relation (16) holds true.

Similar arguments as those used in the proof of relation (16) combined with the fact that since $r^- > \frac{N}{m^-}$ we also have $r^- > \frac{N}{m^+}$ imply that there exists D''_{ϵ} ,

$$\int_{\Omega} |V(x)| \cdot |u|^{m^+} dx \leq \epsilon \int_{\Omega} |\nabla u|^{m^+} dx + D''_{\epsilon} |V|_{r^-} \int_{\Omega} |u|^{m^+} dx, \quad \forall u \in W_0^{1,m^+}(\Omega). \tag{18}$$

Using relation (11) we deduce that $m^- \leq m^+ < (\varphi_1)_0$ and thus, implies that $W_0^{1,(\varphi_1)_0}(\Omega) \subset W_0^{1,m^{\pm}}(\Omega)$. On the other hand, similar arguments as those used in the proof of [21, Lemma 2] show that $W_0^1 L_{\varphi_1}(\Omega) \subset W_0^{1,(\varphi_1)_0}(\Omega)$. The above facts imply that relations (16) and (18) hold true for any $u \in X$. Moreover, in the right-hand sides of inequalities (16) and (18) we can take $|V|_{r(x)}$ instead of $|V|_{r^-}$ since $L^{r(x)}(\Omega)$ is continuously embedded in $L^{r^-}(\Omega)$ via inequality (7).

Finally, we point out that since by (11) we have $(\varphi_2)^0 < m^- \leq m(x) \leq m^+ < (\varphi_1)_0$ for each $x \in \bar{\Omega}$ we deduce that

$$\left| \int_{\Omega} \frac{V(x)}{m(x)} |u|^{m(x)} dx \right| \leq \frac{1}{m^-} \int_{\Omega} |V(x)| \cdot (|u|^{m^-} + |u|^{m^+}) dx, \quad \forall u \in X, \tag{19}$$

and

$$\int_{\Omega} (|\nabla u|^{m^-} + |\nabla u|^{m^+}) dx \leq \int_{\Omega} (|\nabla u|^{(\varphi_2)^0} dx + |\nabla u|^{(\varphi_1)_0}) dx, \quad \forall u \in X. \tag{20}$$

Relations (16), (18), (19), (20), (11) and (15) and [21, Lemma 3] lead to the idea that Lemma 3.1 holds true. \square

Lemma 3.2. *The following relations hold true:*

$$\lim_{\|u\|_1 \rightarrow \infty} \frac{J_V(u)}{I(u)} = \infty, \tag{21}$$

and

$$\lim_{\|u\|_1 \rightarrow 0} \frac{J_V(u)}{I(u)} = \infty. \tag{22}$$

Proof. First, we point out that by (11) $q_2(x) < m^{\pm} < q_1(x)$ for any $x \in \bar{\Omega}$. Thus, it is clear that

$$|u(x)|^{m^-} + |u(x)|^{m^+} \leq 2(|u(x)|^{q_1(x)} + |u(x)|^{q_2(x)}), \quad \forall x \in \bar{\Omega} \text{ and } \forall u \in X.$$

Integrating over Ω the above inequality we infer that

$$\frac{\int_{\Omega} (|u|^{m^-} + |u|^{m^+}) dx}{\int_{\Omega} (|u|^{q_1(x)} + |u|^{q_2(x)}) dx} \leq 2, \quad \forall u \in X. \tag{23}$$

Using Lemma 3.1 we find that for an $\epsilon \in (0, 1)$ there exists $C_{\epsilon} > 0$ such that

$$\frac{J_V(u)}{I(u)} \geq \frac{(1 - \epsilon) \int_{\Omega} (\Phi_1(|\nabla u|) + \Phi_2(|\nabla u|)) dx - C_{\epsilon} |V|_{r(x)} \int_{\Omega} (|u|^{m^-} + |u|^{m^+}) dx}{\frac{1}{q_2} \int_{\Omega} (|u|^{q_1(x)} + |u|^{q_2(x)}) dx},$$

for any $u \in X$.

By the above inequality and relation (23) we deduce that there exist some positive constants $\beta > 0$ and $\gamma > 0$ such that

$$\frac{J_V(u)}{I(u)} \geq \frac{\beta \int_{\Omega} (\Phi_1(|\nabla u|) + \Phi_2(|\nabla u|)) dx}{\int_{\Omega} (|u|^{q_1(x)} + |u|^{q_2(x)}) dx} - \gamma |V|_{r(x)}, \quad \forall u \in X. \tag{24}$$

For any $u \in X$ with $\|u\|_1 > 1$ relation (24) implies:

$$\frac{J_V(u)}{I(u)} \geq \frac{\beta \int_{\Omega} \Phi_1(|\nabla u|) dx}{|u|_{q_1^-}^{q_1^-} + |u|_{q_1^+}^{q_1^+} + |u|_{q_2^-}^{q_2^-} + |u|_{q_2^+}^{q_2^+}} - \gamma |V|_{r(x)}, \quad \forall u \in X \text{ with } \|u\|_1 > 1.$$

Now, taking into account the continuous embedding of X in $L^{q_i^\pm}(\Omega)$ for $i = 1, 2$ (given by relations (11) and (15)) and the result of relation (5) we deduce the existence of a positive constant $\delta > 0$ such that

$$\frac{J_V(u)}{I(u)} \geq \frac{\delta \|u\|_1^{(\varphi_1)_0}}{\|u\|_1^{q_1^-} + \|u\|_1^{q_1^+} + \|u\|_1^{q_2^-} + \|u\|_1^{q_2^+}} - \gamma |V|_{r(x)}, \quad \forall u \in X \text{ with } \|u\|_1 > 1.$$

Since $(\varphi_1)_0 > q_1^+ \geq q_1^- \geq q_2^+ \geq q_2^-$, passing to the limit as $\|u\|_1 \rightarrow \infty$ in the above inequality we deduce that relation (21) holds true.

Relation (15) shows that the space $W_0^1 L_{\Phi_1}(\Omega)$ is continuously embedded in $W_0^1 L_{\Phi_2}(\Omega)$. Thus, if $\|u\|_1 \rightarrow 0$ then $\|u\|_2 \rightarrow 0$.

The above remarks enable us to affirm that for any $u \in X$ with $\|u\|_1 < 1$ small enough we have $\|u\|_2 < 1$.

Using again relation (15) we deduce that $W_0^1 L_{\Phi_2}(\Omega)$ is continuously embedded in $L^{q_i^\pm}(\Omega)$ with $i = 1, 2$. It follows that there exist four positive constants d_{i1} and d_{i2} with $i = 1, 2$ such that

$$\|u\|_2 \geq d_{i1} \cdot |u|_{q_i^+}, \quad \forall u \in W_0^1 L_{\Phi_2}(\Omega) \text{ and } i = 1, 2 \tag{25}$$

and

$$\|u\|_2 \geq d_{i2} \cdot |u|_{q_i^-}, \quad \forall u \in W_0^1 L_{\Phi_2}(\Omega) \text{ and } i = 1, 2. \tag{26}$$

Thus, for any $u \in X$ with $\|u\|_1 < 1$ small enough, relation (24) implies:

$$\frac{J_V(u)}{I(u)} \geq \frac{\beta \int_{\Omega} \Phi_2(|\nabla u|) dx}{|u|_{q_1^-}^{q_1^-} + |u|_{q_1^+}^{q_1^+} + |u|_{q_2^-}^{q_2^-} + |u|_{q_2^+}^{q_2^+}} - \gamma |V|_{r(x)}.$$

Next, relations (4), (25), (26) yield that there exists a constant $\xi > 0$ such that

$$\frac{J_V(u)}{I(u)} \geq \frac{\xi \|u\|_2^{(\varphi_2)_0}}{\|u\|_2^{q_1^-} + \|u\|_2^{q_1^+} + \|u\|_2^{q_2^-} + \|u\|_2^{q_2^+}} - \gamma |V|_{r(x)},$$

for any $u \in X$ with $\|u\|_1 < 1$ small enough. Since $(\varphi_2)_0 < q_2^- \leq q_2^+ \leq q_1^- \leq q_1^+$, passing to the limit as $\|u\|_1 \rightarrow 0$ (and thus, $\|u\|_2 \rightarrow 0$) in the above inequality we deduce that relation (22) holds true. The proof of Lemma 3.2 is complete. \square

Remark 2. We point out that by relation (24) and using similar arguments as in the proof of Theorem 1 (Step 1) in [21] we can find that for V given and satisfying (13) the quotient $\frac{J_V(u)}{I(u)}$ is bounded from below for $u \in X \setminus \{0\}$, i.e. $A(V)$ is a real number. Similarly, it can be proved that $B(V)$ is also a real number.

Lemma 3.3. *There exists $u \in X \setminus \{0\}$ such that $\frac{J_V(u)}{I(u)} = A(V)$.*

Proof. Let $(u_n) \subset X \setminus \{0\}$ be a minimizing sequence for $A(V)$, that is,

$$\lim_{n \rightarrow \infty} \frac{J_V(u_n)}{I(u_n)} = A(V). \tag{27}$$

By relation (21) it is clear that $\{u_n\}$ is bounded in X . Since X is reflexive it follows that there exists $u \in X$ such that, up to a subsequence, (u_n) converges weakly to u in X . On the other hand, similar arguments as those used in the proof of [19, Theorem 2] (see also [21, Step 3]) show that the functional J_0 (obtained for $V = 0$ on Ω) is weakly lower semi-continuous. Thus, we find:

$$\liminf_{n \rightarrow \infty} J_0(u_n) \geq J_0(u). \tag{28}$$

By the compact embedding theorem for Sobolev spaces and assumptions (11), (12) and (13) it follows that X is compactly embedded in $L^{\sigma(x)}(\Omega)$ (where $\sigma(x) = m(x) \cdot r(x)/(r(x) - 1)$) and $L^{q_i(x)}(\Omega)$ with $i = 1, 2$. Thus, (u_n) converges strongly in $L^{\sigma(x)}(\Omega)$ and $L^{q_i(x)}(\Omega)$ with $i = 1, 2$. Then, by relations (7) and (15) it follows that

$$\lim_{n \rightarrow \infty} I(u_n) = I(u), \tag{29}$$

and

$$\lim_{n \rightarrow \infty} \int_{\Omega} V(x)|u_n|^{m(x)} dx = \int_{\Omega} V(x)|u|^{m(x)} dx. \tag{30}$$

Relations (28), (29) and (30) imply that if $u \neq 0$, then

$$\frac{J_V(u)}{I(u)} = A(V).$$

Thus, in order to conclude that the lemma holds true it is enough to show that u is not trivial. Assume by contradiction the contrary. Then u_n converges weakly to 0 in X and strongly in $L^{s(x)}(\Omega)$ for any $s(x) \in C(\overline{\Omega})$ with $1 < s(x) < \frac{N(\varphi_1)_0}{N - (\varphi_1)_0}$ on $\overline{\Omega}$. In other words, we will have:

$$\lim_{n \rightarrow \infty} I(u_n) = 0, \tag{31}$$

and

$$\lim_{n \rightarrow \infty} \int_{\Omega} V(x)|u_n|^{m(x)} dx = 0. \tag{32}$$

Letting $\epsilon \in (0, |A(V)|)$ be fixed by relation (27) we deduce that for n large enough we have:

$$|J_V(u_n) - A(V)I(u_n)| < \epsilon I(u_n),$$

or

$$(|A(V)| - \epsilon)I(u_n) < J_V(u_n) < (|A(V)| + \epsilon)I(u_n).$$

Passing to the limit in the above inequalities and taking into account that relation (31) holds true we find:

$$\lim_{n \rightarrow \infty} J_V(u_n) = 0.$$

Next, by relation (32) we get:

$$\lim_{n \rightarrow \infty} J_0(u_n) = 0.$$

That fact combined with relation (4) implies that actually u_n converges strongly to 0 in X , i.e. $\lim_{n \rightarrow \infty} \|u_n\|_1 = 0$. By this information and relation (22) we get:

$$\lim_{n \rightarrow \infty} \frac{J_V(u_n)}{I(u_n)} = \infty,$$

and this is a contradiction. Thus, $u \neq 0$. The proof of Lemma 3.3 is complete. \square

By Lemma 3.3 we conclude that there exists $u \in X \setminus \{0\}$ such that

$$\frac{J_V(u)}{I(u)} = A(V) = \inf_{w \in X \setminus \{0\}} \frac{J_V(w)}{I(w)}. \tag{33}$$

Then, for any $w \in X$ we have:

$$\left. \frac{d}{d\epsilon} \frac{J_V(u + \epsilon w)}{I(u + \epsilon w)} \right|_{\epsilon=0} = 0.$$

A simple computation yields

$$\langle J'_V(u), w \rangle I(u) - J_V(u) \langle I'(u), w \rangle = 0, \tag{34}$$

for all $w \in X$. Relation (34) combined with the fact that $J_V(u) = A(V) \cdot I(u)$ and $I(u) \neq 0$ implies the fact that $A(V)$ is an eigenvalue of problem (1).

Next, we show that any $\lambda \in (A(V), \infty)$ is an eigenvalue of problem (1).

Let $\lambda \in (A(V), \infty)$ be arbitrary but fixed. Define $T_{V,\lambda} : X \rightarrow \mathbb{R}$ by:

$$T_{V,\lambda}(u) = J_V(u) - \lambda I(u).$$

Clearly, $T_{V,\lambda} \in C^1(X, \mathbb{R})$ with

$$\langle T'_{V,\lambda}(u), v \rangle = \langle J'_V(u), v \rangle - \lambda \langle I'(u), v \rangle, \quad \forall u \in X.$$

Thus, λ is an eigenvalue of problem (1) if and only if there exists $u_\lambda \in X \setminus \{0\}$ a critical point of $T_{V,\lambda}$.

With similar arguments as in the proof of relation (21) we can show that $T_{V,\lambda}$ is coercive, i.e. $\lim_{\|u\| \rightarrow \infty} T_{V,\lambda}(u) = \infty$. On the other hand, as we have already remarked, similar arguments as those used in the proof of [19, Theorem 2] show that the functional $T_{V,\lambda}$ is weakly lower semi-continuous. These two facts enable us to apply [27, Theorem 1.2] in order to prove that there exists $u_\lambda \in X$ a global minimum point of $T_{V,\lambda}$ and thus, a critical point of $T_{V,\lambda}$. It is enough to show that u_λ is not trivial. Indeed, since $A(V) = \inf_{u \in X \setminus \{0\}} \frac{J_V(u)}{I(u)}$ and $\lambda > A(V)$ it follows that there exists $v_\lambda \in X$ such that

$$J_V(v_\lambda) < \lambda I(v_\lambda),$$

or

$$T_{V,\lambda}(v_\lambda) < 0.$$

Thus,

$$\inf_X T_{V,\lambda} < 0$$

and we conclude that u_λ is a nontrivial critical point of $T_{V,\lambda}$, or λ is an eigenvalue of problem (1).

Finally, we prove that each $\lambda < B(V)$ is not an eigenvalue of problem (1). With that end in view we assume by contradiction that there exists $\lambda < B(V)$ an eigenvalue of problem (1). It follows that there exists $u_\lambda \in X \setminus \{0\}$ such that

$$\langle J'_V(u_\lambda), u_\lambda \rangle = \lambda \langle I'(u_\lambda), u_\lambda \rangle.$$

Since $u_\lambda \neq 0$ we have $\langle I'(u_\lambda), u_\lambda \rangle > 0$. Using that fact and the definition of $B(V)$ it follows that the following relation holds true:

$$\langle J'_V(u_\lambda), u_\lambda \rangle = \lambda \langle I'(u_\lambda), u_\lambda \rangle < B(V) \langle I'(u_\lambda), u_\lambda \rangle \leq \langle J'_V(u_\lambda), u_\lambda \rangle.$$

Obviously, this is a contradiction. We deduce that each $\lambda \in (-\infty, B(V))$ is not an eigenvalue of problem (1). Furthermore, it is clear that $A(V) \geq B(V)$.

The proof of Theorem 2.1 is complete.

Remark 3. We point out that in the case when $V = 0$ in Ω the same arguments as in the proof of Theorem 1 (Step 1) in [21] assure that $A(0) > 0$.

4. Proof of Theorem 2.2

Let S be a convex, bounded and closed subset of $L^{r(x)}(\Omega)$, and

$$A_\star := \inf_{V \in S} A(V).$$

Clearly, relation (24) assures that A_\star is finite.

On the other hand, let $(V_n) \subset S$ be a minimizing sequence for A_\star , i.e.

$$A(V_n) \rightarrow A_\star, \quad \text{as } n \rightarrow \infty.$$

Obviously, (V_n) is a bounded sequence and thus, there exists $V_\star \in L^{r(x)}(\Omega)$ such that V_n converges weakly to V_\star in $L^{r(x)}(\Omega)$. Moreover, since S is convex and closed it is also weakly closed (see, e.g., Brezis [5, Theorem III.7]) and consequently $V_\star \in S$.

Next, we will show that $A(V_\star) = A_\star$.

Indeed, by Theorem 2.1 we deduce that for each positive integer n there exists $u_n \in X \setminus \{0\}$ such that

$$\frac{J_{V_n}(u_n)}{I(u_n)} = A(V_n). \tag{35}$$

Since $(A(V_n))$ is a bounded sequence and by relation (24) we have:

$$\frac{J_{V_n}(u_n)}{I(u_n)} \geq \beta \frac{J_0(u_n)}{I(u_n)} - C, \quad \text{for any } n,$$

where C is a positive constant, we infer that (u_n) is bounded in X and it cannot contain a subsequence converging to 0 (otherwise we obtain a contradiction by applying Lemma 3.2). Thus, there exists $u_0 \in X \setminus \{0\}$ such that (u_n) converges weakly to u_0 in X . Using relation (12) (and thus, $W_0^1 L_{\Phi_1}(\Omega) \subset W_0^{1,(\varphi_1)_0}(\Omega)$) and the Rellich–Kondrachov theorem we deduce that (u_n) converges strongly to u_0 in $L^{s(x)}(\Omega)$ for any $s(x) \in C(\overline{\Omega})$ satisfying $1 < s(x) < \frac{N(\varphi_1)_0}{N - (\varphi_1)_0}$ for any $x \in \overline{\Omega}$. In particular, using conditions (11), (12) and (13) we get that (u_n) converges to u_0 in $L^{m(x)}(\Omega)$ and in $L^{m(x) \cdot r'(x)}(\Omega)$ where $r'(x) = \frac{r(x)}{r(x)-1}$. Using that information, inequality (7) and the fact that $V_\star \in L^{r(x)}(\Omega)$ and (V_n) is bounded in $L^{r(x)}(\Omega)$ we find:

$$\lim_{n \rightarrow \infty} \int_{\Omega} \frac{V_\star(x)}{m(x)} |u_n|^{m(x)} dx = \int_{\Omega} \frac{V_\star(x)}{m(x)} |u_0|^{m(x)} dx, \tag{36}$$

and

$$\lim_{n \rightarrow \infty} \int_{\Omega} \left(\frac{V_n(x)}{m(x)} |u_n|^{m(x)} - \frac{V_n(x)}{m(x)} |u_0|^{m(x)} \right) dx = 0. \tag{37}$$

On the other hand, since (V_n) converges weakly to V_\star in $L^{r(x)}(\Omega)$ and $u_0 \in L^{m(x) \cdot r'(x)}(\Omega)$, where $r'(x) = \frac{r(x)}{r(x)-1}$, we deduce:

$$\lim_{n \rightarrow \infty} \int_{\Omega} \frac{V_n(x)}{m(x)} |u_0|^{m(x)} dx = \int_{\Omega} \frac{V_\star(x)}{m(x)} |u_0|^{m(x)} dx. \tag{38}$$

Combining the equality,

$$\begin{aligned} & \int_{\Omega} \frac{V_\star(x)}{m(x)} |u_n|^{m(x)} dx - \int_{\Omega} \frac{V_n(x)}{m(x)} |u_n|^{m(x)} dx \\ &= \int_{\Omega} \frac{V_\star(x)}{m(x)} |u_n|^{m(x)} dx - \int_{\Omega} \frac{V_\star(x)}{m(x)} |u_0|^{m(x)} dx + \int_{\Omega} \frac{V_\star(x)}{m(x)} |u_0|^{m(x)} dx - \int_{\Omega} \frac{V_n(x)}{m(x)} |u_0|^{m(x)} dx \\ & \quad + \int_{\Omega} \frac{V_n(x)}{m(x)} |u_0|^{m(x)} dx - \int_{\Omega} \frac{V_n(x)}{m(x)} |u_n|^{m(x)} dx, \end{aligned}$$

with relations (36), (37) and (38) we get:

$$\lim_{n \rightarrow \infty} \int_{\Omega} \left(\frac{V_{\star}(x)}{m(x)} |u_n|^{m(x)} - \frac{V_n(x)}{m(x)} |u_n|^{m(x)} \right) dx = 0. \tag{39}$$

Since

$$A(V_{\star}) = \inf_{u \in X \setminus \{0\}} \frac{J_{V_{\star}}(u)}{I(u)},$$

it follows that

$$A(V_{\star}) \leq \frac{J_{V_{\star}}(u_n)}{I(u_n)}.$$

Combining the above inequality and equality (35) we obtain:

$$A(V_{\star}) \leq \frac{J_{V_{\star}}(u_n) - J_{V_n}(u_n)}{I(u_n)} + A(V_n).$$

Taking into account the result of relation (39), the fact that $I(u_n)$ is bounded and does not converge to 0 and $(A(V_n))$ converges to A_{\star} then passing to the limit as $n \rightarrow \infty$ in the last inequality we infer that

$$A(V_{\star}) \leq A_{\star}.$$

But using the definition of A_{\star} and the fact that $V_{\star} \in S$ we conclude that actually

$$A(V_{\star}) = A_{\star}.$$

The proof of Theorem 2.2 is complete.

5. Proof of Theorem 2.3

(a) First, we show that function A_{\star} is not constant. Indeed, by Remark 3 we point out that $A_{\star}(0) = A(0) > 0$. On the other hand, by [21, Theorem 1] it follows that

$$\lambda_m := \inf_{u \in X \setminus \{0\}} \frac{\int_{\Omega} \Phi_1(|\nabla u|) dx + \int_{\Omega} \Phi_2(|\nabla u|) dx}{\int_{\Omega} \frac{1}{m(x)} |u|^{m(x)} dx} > 0.$$

Moreover, [21, Lemma 5] implies that there exists $u_m \in X \setminus \{0\}$ such that

$$\lambda_m = \frac{\int_{\Omega} \Phi_1(|\nabla u_m|) dx + \int_{\Omega} \Phi_2(|\nabla u_m|) dx}{\int_{\Omega} \frac{1}{m(x)} |u_m|^{m(x)} dx}.$$

Thus, taking $V_m(x) = -\lambda_m$ for all $x \in \Omega$ it is clear that $V_m \in L^{\infty}(\Omega) \subset L^{r(x)}(\Omega)$, and

$$\frac{J_{V_m}(u_m)}{I(u_m)} = 0.$$

It follows that

$$A(V_m) \leq 0,$$

and we find,

$$A_{\star}(\lambda_m) \leq 0.$$

We conclude that A_{\star} is not constant. Furthermore, we point out that a similar proof as those presented above can show that function A_{\star} takes also negative values. To support that idea we just notice that by [21, Theorem 1, Step 3] for each $\lambda > \lambda_m$ there exists $u_{\lambda} \in X \setminus \{0\}$ such that taking $V_{\lambda} = -\lambda$ for all $x \in \Omega$ we have:

$$\frac{J_{V_{\lambda}}(u_{\lambda})}{I(u_{\lambda})} < 0.$$

Next, we point out that A_\star decreases monotonically. Indeed, if we consider $0 \leq R_1 < R_2$ then it is clear that $\bar{B}_{R_1}(0) \subset \bar{B}_{R_2}(0)$. Then the definition of function A_\star implies $A_\star(R_1) \geq A_\star(R_2)$.

(b) Finally, we show that the function A_\star is continuous. Let $R > 0$ and $t \in (0, R)$ be fixed. We will verify that $\lim_{t \searrow 0} A_\star(R+t) = \lim_{t \searrow 0} A_\star(R-t) = A_\star(R)$.

First, we prove that $\lim_{t \searrow 0} A_\star(R+t) = A_\star(R)$. By Theorem 2.3(a) we have:

$$A_\star(R) \geq A_\star(R+t).$$

Moreover, by Theorem 2.2 it follows that there exists $V_{R+t} \in \bar{B}_{R+t}(0)$ (i.e. $|V_{R+t}|_{r(x)} \leq R+t$) such that

$$A(V_{R+t}) = A_\star(R+t).$$

Taking now $V_{R,t} := \frac{R}{R+t} V_{R+t}$ we have:

$$|V_{R,t}|_{r(x)} = \frac{R}{R+t} |V_{R+t}|_{r(x)} \leq R,$$

or $V_{R,t} \in \bar{B}_R(0)$. Therefore, obviously, we have $A(V_{R,t}) \geq A_\star(R)$.

On the other hand, by Theorem 2.1 there exists $u_t \in X \setminus \{0\}$ such that

$$A(V_{R+t}) = \frac{J_{V_{R+t}}(u_t)}{I(u_t)}.$$

Combining the above pieces of information we find:

$$\begin{aligned} A_\star(R+t) &= A(V_{R+t}) = \frac{J_{V_{R+t}}(u_t)}{I(u_t)} = \frac{J_{\frac{R+t}{R} V_{R,t}}(u_t)}{I(u_t)} = \frac{R+t}{R} \cdot \frac{J_{V_{R,t}}(u_t)}{I(u_t)} - \frac{t}{R} \cdot \frac{J_0(u_t)}{I(u_t)} \\ &\geq \frac{R+t}{R} \cdot A_\star(R) - \frac{t}{R} \cdot \frac{J_0(u_t)}{I(u_t)}. \end{aligned}$$

On the other hand, by relation (24) we have that for each $t \in (0, R)$ it holds:

$$\begin{aligned} A_\star(R) &\geq A_\star(R+t) = A(V_{R+t}) = \frac{J_{V_{R+t}}(u_t)}{I(u_t)} \geq \beta_1 \cdot \frac{J_0(u_t)}{I(u_t)} - \gamma \cdot |V_{R+t}|_{r(x)} \\ &= \beta_1 \cdot \frac{J_0(u_t)}{I(u_t)} - \gamma \cdot 2R, \end{aligned}$$

where $\beta_1 > 0$ and $\gamma > 0$ are real constants.

Combining the last two inequalities we deduce that

$$A_\star(R) \geq A_\star(R+t) \geq \frac{R+t}{R} \cdot A_\star(R) - \frac{t}{R} \cdot \frac{A_\star(R) + \gamma \cdot 2R}{\beta_1},$$

for each $t \in (0, R)$.

We conclude that

$$\lim_{t \searrow 0} A_\star(R+t) = A_\star(R).$$

In the following we argue that $\lim_{t \searrow 0} A_\star(R-t) = A_\star(R)$.

Obviously,

$$A_\star(R) \leq A_\star(R-t), \quad \forall t \in (0, R).$$

By Theorem 2.2 there exists $V_R \in \bar{B}_R(0)$ such that

$$A_\star(R) = A(V_R).$$

Moreover, by Theorem 2.1 there exists $u_0 \in X \setminus \{0\}$ such that

$$A(V_R) = \frac{J_{V_R}(u_0)}{I(u_0)}.$$

Define now:

$$V_t := \frac{R-t}{R} V_R, \quad \forall t \in (0, R).$$

Clearly, $V_t \in \overline{B}_{R-t}(0)$. Thus, it is clear that

$$\frac{J_{V_t}(u_0)}{I(u_0)} \geq A_\star(R-t), \quad \forall t \in (0, R).$$

Taking into account the above information we find:

$$\begin{aligned} A_\star(R) &= A(V_R) = \frac{J_{V_R}(u_0)}{I(u_0)} = \frac{J_{\frac{R}{R-t}V_t}(u_0)}{I(u_0)} = \frac{J_{V_t}(u_0)}{I(u_0)} + \frac{t}{R-t} \cdot \frac{\int_{\Omega} \frac{V_t(x)}{m(x)} |u_0|^{m(x)} dx}{I(u_0)} \\ &\geq A_\star(R-t) + \frac{t}{R} \cdot \frac{\int_{\Omega} \frac{V_R(x)}{m(x)} |u_0|^{m(x)} dx}{I(u_0)}, \quad \forall t \in (0, R). \end{aligned}$$

We infer

$$\lim_{t \searrow 0} A_\star(R-t) = A_\star(R).$$

It follows that function A_\star is continuous. The proof of Theorem 2.3 is complete.

Remark 4. By Theorem 2.3(a) we get that A_\star decreases monotonically. We notice that in the particular case when $q_1(x) = m(x) = q_2(x) = q$ for each $x \in \overline{\Omega}$, where $q > 1$ is a real number for which conditions (11), (12) and (13) are fulfilled, the above quoted result can be improved, in the sense that we can show that, actually, function A_\star is strictly decreasing on $[0, \infty)$. Indeed, letting $0 \leq R_1 < R_2$ be given, by Theorem 2.2 we deduce that there exists $V_1 \in \overline{B}_{R_1}(0)$ such that

$$A(V_1) = A_\star(R_1).$$

Then for each real number $t \in (0, R_2 - R_1)$ we have $V_1 - t \in \overline{B}_{R_2}(0)$ since $|V_1 - t|_{r(x)} \leq |V_1|_{r(x)} + t \leq R_2$. Next, by Theorem 2.1 there exists $u_1 \in X \setminus \{0\}$ such that

$$A(V_1) = \frac{J_{V_1}(u_1)}{I(u_1)}.$$

Taking into account all the above remarks we infer:

$$A_\star(R_1) - \frac{t}{2} = A(V_1) - \frac{t}{2} = \frac{J_{V_1}(u_1)}{I(u_1)} - \frac{t}{2} = \frac{J_{V_1-t}(u_1)}{I(u_1)} \geq A(V_1 - t) \geq A_\star(R_2),$$

or

$$A_\star(R_1) > A_\star(R_2).$$

In the end of this remark we consider that it is important to highlight the idea that the above proof supports the fact that in the case when we manipulate homogeneous quantities we obtain better results than in the case when we deal with non-homogeneous quantities.

Remark 5. We point out that by Theorem 2.3(b) we deduce that

$$A_\star(R) = \inf_{s \leq R} A_\star(s) \quad \text{and} \quad A_\star(R) = \sup_{s \geq R} A_\star(s).$$

Remark 6. We also point out that function A_\star can be used in order to define a *continuous set function* on a subset of $L^{r(x)}(\Omega)$. We still denote each closed ball centered in the origin of radius R from $L^{r(x)}(\Omega)$ by $\overline{B}_R(0)$, i.e.

$$\overline{B}_R(0) := \{u \in L^{r(x)}(\Omega); |u|_{r(x)} \leq R\}.$$

By Theorem 2.3(b) we deduce that A_\star is a continuous function. By the proof of Theorem 2.3(a) we have $A_\star(0) > 0$ and there exists $R_1 > 0$ such that $A_\star(R_1) < 0$. Thus, we infer that there exists $R_0 > 0$ such that $A_\star(R_0) = 0$.

We define:

$$\Gamma = \{\bar{B}_R(0) \setminus \bar{B}_{R_0}(0); R \geq R_0\} \subset L^{r(x)}(\Omega),$$

and $\mu: \Gamma \rightarrow [0, \infty)$ by:

$$\mu(\bar{B}_R(0) \setminus \bar{B}_{R_0}(0)) = -A_*(R), \quad \forall R \geq R_0.$$

By Theorem 2.3(a) we find that function μ has the following properties:

- 1) $\mu(\emptyset) = 0$;
- 2) For each $S_1, S_2 \in \Gamma$ such that $S_1 \subset S_2$ we have $\mu(S_1) \leq \mu(S_2)$.

Thus, μ is a *set function* on Γ . By Theorem 2.3(b) and Remark 4 we have that for each $S \subset \Gamma$ it holds true that

$$\mu(S) = \sup_{T \subset S} \mu(T) \quad \text{and} \quad \mu(S) = \inf_{T \supseteq S} \mu(T).$$

We conclude that μ is a *continuous set function* on Γ .

Acknowledgements

The first two named authors have been supported by Grant CNCSIS PNII-79/2007 “Procese Neliniare Degenerate și Singulare”. V. Rădulescu also acknowledges support through Grant CNCSIS PCCE-55/2008 “Sisteme Diferențiale în Analiza Neliniară și Aplicații”. D. Repovš and V. Rădulescu were supported by the Slovenian Research Agency grants 1000-08-780004, P1-0292-0101 and J1-9643-0101.

References

- [1] M.S. Ashbaugh, E.M. Harrell, Maximal and minimal eigenvalues and their associated nonlinear equations, *J. Math. Phys.* 28 (1987) 1770–1786.
- [2] D.R. Adams, L.I. Hedberg, *Function Spaces and Potential Theory*, Grundlehren der Mathematischen Wissenschaften (Fundamental Principles of Mathematical Sciences), vol. 314, Springer-Verlag, Berlin, 1996.
- [3] R. Adams, *Sobolev Spaces*, Academic Press, New York, 1975.
- [4] J.F. Bonder, L.M. Del Pezzo, An optimization problem for the first eigenvalue of the p -Laplacian plus a potential, *Comm. Pure Appl. Anal.* 5 (2006) 675–690.
- [5] H. Brezis, *Analyse fonctionnelle: théorie, méthodes et applications*, Masson, Paris, 1992.
- [6] Ph. Clément, M. García-Huidobro, R. Manásevich, K. Schmitt, Mountain pass type solutions for quasilinear elliptic equations, *Calc. Var.* 11 (2000) 33–62.
- [7] Ph. Clément, B. de Pagter, G. Sweers, F. de Thélin, Existence of solutions to a semilinear elliptic system through Orlicz–Sobolev spaces, *Mediterr. J. Math.* 1 (2004) 241–267.
- [8] D.E. Edmunds, J. Lang, A. Nekvinda, On $L^{p(x)}$ norms, *Proc. Roy. Soc. London Ser. A* 455 (1999) 219–225.
- [9] D.E. Edmunds, J. Rákosník, Density of smooth functions in $W^{k,p(x)}(\Omega)$, *Proc. Roy. Soc. London Ser. A* 437 (1992) 229–236.
- [10] D.E. Edmunds, J. Rákosník, Sobolev embedding with variable exponent, *Studia Math.* 143 (2000) 267–293.
- [11] H. Egnell, Extremal properties of the first eigenvalue of a class of elliptic eigenvalue problems, *Ann. Scuola Norm. Sup. Pisa* 14 (1987) 1–48.
- [12] X. Fan, Remarks on eigenvalue problems involving the $p(x)$ -Laplacian, *J. Math. Anal. Appl.* 352 (2009) 85–98.
- [13] X. Fan, Q. Zhang, D. Zhao, Eigenvalues of $p(x)$ -Laplacian Dirichlet problem, *J. Math. Anal. Appl.* 302 (2005) 306–317.
- [14] M. García-Huidobro, V.K. Le, R. Manásevich, K. Schmitt, On principal eigenvalues for quasilinear elliptic differential operators: An Orlicz–Sobolev space setting, *Nonlinear Differential Equations Appl. (NoDEA)* 6 (1999) 207–225.
- [15] J.P. Gossez, Nonlinear elliptic boundary value problems for equations with rapidly (or slowly) increasing coefficients, *Trans. Amer. Math. Soc.* 190 (1974) 163–205.
- [16] O. Kováčik, J. Rákosník, On spaces $L^{p(x)}$ and $W^{1,p(x)}$, *Czechoslovak Math. J.* 41 (1991) 592–618.
- [17] M. Mihăilescu, V. Rădulescu, A multiplicity result for a nonlinear degenerate problem arising in the theory of electrorheological fluids, *Proc. Roy. Soc. A: Math., Phys. Engrg. Sci.* 462 (2006) 2625–2641.
- [18] M. Mihăilescu, V. Rădulescu, On a nonhomogeneous quasilinear eigenvalue problem in Sobolev spaces with variable exponent, *Proc. Amer. Math. Soc.* 135 (2007) 2929–2937.
- [19] M. Mihăilescu, V. Rădulescu, Existence and multiplicity of solutions for quasilinear nonhomogeneous problems: an Orlicz–Sobolev space setting, *J. Math. Anal. Appl.* 330 (2007) 416–432.
- [20] M. Mihăilescu, V. Rădulescu, Continuous spectrum for a class of nonhomogeneous differential operators, *Manuscripta Math.* 125 (2008) 157–167.

- [21] M. Mihăilescu, V. Rădulescu, Eigenvalue problems associated to nonhomogeneous differential operators in Orlicz–Sobolev spaces, *Anal. Appl.* 6 (1) (2008) 1–16.
- [22] M. Mihăilescu, V. Rădulescu, Neumann problems associated to nonhomogeneous differential operators in Orlicz–Sobolev spaces, *Ann. Inst. Fourier* 58 (6) (2008) 2087–2111.
- [23] M. Mihăilescu, V. Rădulescu, Spectrum in an unbounded interval for a class of nonhomogeneous differential operators, *Bull. London Math. Soc.* 40 (6) (2008) 972–984.
- [24] J. Musielak, *Orlicz Spaces and Modular Spaces*, Lecture Notes in Mathematics, vol. 1034, Springer-Verlag, Berlin, 1983.
- [25] M.M. Rao, Z.D. Ren, *Theory of Orlicz Spaces*, Marcel Dekker, Inc., New York, 1991.
- [26] S. Samko, B. Vakulov, Weighted Sobolev theorem with variable exponent for spatial and spherical potential operators, *J. Math. Anal. Appl.* 310 (2005) 229–246.
- [27] M. Struwe, *Variational Methods: Applications to Nonlinear Partial Differential Equations and Hamiltonian Systems*, Springer, Heidelberg, 1996.